

Incomparable, non isomorphic and minimal Banach spaces

Christian Rosendal

March 2004

Abstract

A Banach space contains either a minimal subspace or a continuum of incomparable subspaces. General structure results for analytic equivalence relations are applied in the context of Banach spaces to show that if E_0 does not reduce to isomorphism of the subspaces of a space, in particular, if the subspaces of the space admit a classification up to isomorphism by real numbers, then any subspace with an unconditional basis is isomorphic to its square and hyperplanes and has an isomorphically homogeneous subsequence.

1 Introduction.

This paper contains results in the intersection of the geometry of Banach spaces and descriptive set theory. The general problem of our study is a generalisation of the homogeneous space problem. Namely, what can be said about a Banach space with “few” non isomorphic subspaces? In particular, will such a space necessarily satisfy more regularity properties than a general space? Will it necessarily have subspaces of a given type?

The paper is divided into two parts, of which the first contains a proof of the following:

Theorem 1 *Let X be an infinite dimensional Banach space. Then X contains either a minimal subspace or a continuum of pairwise incomparable subspaces.*

Recall that two spaces are said to be incomparable if neither of them embed into the other, and a space is minimal if it embeds into all of its infinite dimensional subspaces.

The homogeneous space problem, which was solved in the positive by the combined efforts of Gowers [6], Komorowski and Tomczak-Jaegermann [13], is the problem of whether any infinite dimensional space, isomorphic to all its infinite dimensional subspaces, must necessarily be isomorphic to ℓ_2 . As a continuation of this one can ask how many isomorphism classes of subspaces a non Hilbertian space has to contain. Infinitely many? A continuum? Even for some of the classical spaces this question is still open, though recent progress has been made by Ferenczi and Galego [3].

Our theorem and proof turn out to have something to say about the following two problems of Gowers. ([6], Problems 7.9 and 7.10):

- Determine which partial orders that can be realised as the set of subspaces of an infinite dimensional Banach space under the relation of embeddability. Or at least find strong conditions such a partial order must necessarily satisfy.
- Find further applications of the main determinacy result in [6]. In particular, are there any applications that need its full strength, i.e., that need it to hold for analytic and not just open sets?

Our Theorem 1 says that any such partial order must either have a minimal element or an antichain of continuum size. And, as will be evident, the proof does in fact very much need the full strength of the determinacy result.

We mention that our proof relies heavily on methods of logic and we have therefore included a short review of the most basic notions of set theory indispensable to understand the proof. Also for the benefit of the non analyst we recall some standard notions from Banach space theory.

Before presenting the results of the second part we will first need this brief review.

1.1 Descriptive set theory.

Our general reference for descriptive set theory will be the book by Kechris, [12], whose notation will be adopted here.

A *Polish* space is a separable completely metrisable space. A measurable space, whose algebra of measurable sets are the Borel sets of some Polish

topology, is said to be *standard Borel*. These spaces turn out to be completely classified up to Borel isomorphism by their cardinality, that can either be countable or equal to that of the continuum. A subset of a standard Borel space is *analytic* if it is the image by a Borel function of some standard Borel space and *coanalytic* if its complement is so. It is *C-measurable* if it belongs to the smallest σ -algebra containing the Borel sets and closed under the Souslin operation. In particular, analytic sets are *C-measurable* as they can be obtained by the Souslin operation applied to a sequence of Borel sets. *C-measurable* sets in Polish spaces satisfy most of the classical regularity properties, such as universal measurability and the Baire property. We denote by Σ_1^1 , Π_1^1 and Σ_2^1 the classes of analytic, coanalytic and Borel images of coanalytic sets respectively. A classic result of Sierpinski states that any Σ_2^1 set is the union of $\aleph_1$ Borel sets.

Let X be a Polish space and $\mathcal{F}(X)$ denote the set of closed subsets of X . We endow $\mathcal{F}(X)$ with the following σ -algebra that renders it a standard Borel space. The generators are the following sets, where U varies over the open subsets of X :

$$\{F \in \mathcal{F}(X) \mid F \cap U \neq \emptyset\}$$

The resulting measurable space is called the Effros Borel space of X .

Fix some basis $\{U_n\}$ for the space $C(2^{\mathbb{N}})$ and define the Borel set $\mathfrak{B}$ by:

$$\begin{aligned} \mathfrak{B} = \{F \in \mathcal{F}(C(2^{\mathbb{N}})) \mid \forall n (0 \in U_n \rightarrow F \cap U_n \neq \emptyset) \wedge \forall n, m, l \forall r, t \in \mathbb{Q} \\ (rU_n + tU_m \subseteq U_l \wedge F \cap U_n \neq \emptyset \wedge F \cap U_m \neq \emptyset \rightarrow F \cap U_l \neq \emptyset)\} \end{aligned}$$

This evidently consists of all the closed linear subspaces of $C(2^{\mathbb{N}})$ and, as $C(2^{\mathbb{N}})$ is isometrically universal for separable Banach spaces, any separable Banach space has an isometric copy in $\mathfrak{B}$. We can therefore view $\mathfrak{B}$ as the standard Borel space of all separable Banach spaces. When one wants to restrict the attention to the subspaces of some particular space X one only needs to consider the Borel subset $\{Y \in \mathfrak{B} \mid Y \subseteq X\}$. Moreover, it is not hard to see that most reasonably definable properties and relations are Σ_2^1 in $\mathfrak{B}$ or $\mathfrak{B}^n$; for example, the relations of isometry and isomorphism are both analytic in $\mathfrak{B}^2$ exactly as expected.

A theme of descriptive set theory, that has been extensively developed the last fifteen years or so, is the Borel reducibility ordering of analytic equivalence relations on standard Borel spaces.

This ordering is defined as follows: Suppose $E \subset X^2$ and $F \subset Y^2$ are analytic equivalence relations on standard Borel spaces X and Y . We say

that E is *Borel reducible* to F , in symbols $E \leq_B F$, if there is a Borel measurable function $f : X \rightarrow Y$ such that for all $x, y \in X$:

$$xEy \iff f(x)Ff(y)$$

Moreover, when X and Y are Polish and f can be taken to be continuous, we write $E \leq_c F$.

Heuristically, X represents a class of mathematical objects (e.g., separable Banach spaces) that we wish to classify up to E -equivalence (e.g., isomorphism) by complete invariants belonging to some other category of mathematical objects. A reduction $f : X \rightarrow Y$ of E to F corresponds then to a classification of X -objects up to E -equivalence by Y -objects up to F -equivalence.

Another way of viewing the Borel reducibility ordering is as a refinement of the concept of cardinality. It provides a concept of relative cardinality for quotient spaces in the absence of the axiom of choice. For a reduction of E to F is essentially an injection of X/E into Y/F admitting a Borel lifting from X to Y .

A few words on the power of the continuum: We say that an analytic equivalence relation E on a standard Borel space X has a *continuum of classes* if there is an uncountable Borel set $B \subset X$ consisting of pairwise E -inequivalent points. This is known to be stronger than just demanding that there should be some bijection between the set of classes and $\mathbb{R}$. There are for example analytic equivalence relations having exactly $\aleph_1$ many classes, but not having a continuum of classes (in the above sense) in any model of set theory. But an uncountable Borel set is always Borel isomorphic to $\mathbb{R}$, independently of the size of the continuum.

If A is some infinite subset of $\mathbb{N}$, we denote by $[A]^\mathbb{N}$ the space of all infinite subsets of A equipped with the topology induced by the product topology on 2^A . Furthermore, for two sets A and B we write $A \subset^* B$ iff $A \setminus B$ is finite. Then $A \subsetneq^* B$ iff $A \subset^* B$ but $B \not\subset^* A$. Also, when $A \subset \mathbb{N}$ and $k \in \mathbb{N}$ we let $A/k = \{n \in A \mid n > k\}$. We will occasionally also consider natural numbers as ordinals, so that $n = \{0, 1, \dots, n-1\}$.

We will repeatedly use the following result of Ellentuck extending results of Galvin-Prikry for Borel sets and Silver for analytic sets: if $\mathcal{A} \subset [\mathbb{N}]^\mathbb{N}$ is a C -measurable set, then there is some $A \in [\mathbb{N}]^\mathbb{N}$ with either $[A]^\mathbb{N} \subset \mathcal{A}$ or $[A]^\mathbb{N} \cap \mathcal{A} = \emptyset$.

This has the consequence that if $f : [\mathbb{N}]^\mathbb{N} \rightarrow X$ is some C -measurable

function with values in some Polish space X , then there is some $A \in [\mathbb{N}]^{\mathbb{N}}$ such that f 's restriction to $[A]^{\mathbb{N}}$ is continuous.

Among the simpler analytic equivalence relations are those that admit a classification by real numbers, i.e., those that are Borel reducible to the identity relation on $\mathbb{R}$. These are said to be *smooth*. It turns out that among Borel equivalence relations there is a minimum, with respect to $\leq_B$, non smooth one, which we denote by E_0 (see [8]). It is defined on $[\mathbb{N}]^{\mathbb{N}}$ as the relation of eventual agreement, i.e.:

$$AE_0B \iff \exists n \ A/n = B/n$$

To see that E_0 is non smooth, suppose towards a contradiction that $f: [\mathbb{N}]^{\mathbb{N}} \rightarrow \mathbb{R}$ is a Borel function such that $AE_0B \iff f(A) = f(B)$. Then there is some infinite $C \subset \mathbb{N}$ such that the restriction of f to $[C]^{\mathbb{N}}$ is continuous. But, as the equivalence class of C is dense in $[C]^{\mathbb{N}}$, this means that f is constant on $[C]^{\mathbb{N}}$, contradicting that $[C]^{\mathbb{N}}$ intersects more than one equivalence class.

On the other hand any uncountable Borel set $B \subset [\mathbb{N}]^{\mathbb{N}}$ of pairwise almost disjoint sets will witness that E_0 has a continuum of classes.

From this it follows that any analytic equivalence relation to which E_0 reduces has a continuum of classes, but does not admit a classification by real numbers.

After these preliminary remarks we can state our second result.

Theorem 2 *Let X be a Banach space with an unconditional basis (e_n) . If E_0 does not Borel reduce to isomorphism between subspaces generated by subsequences of the basis (and in particular if these admit a classification by real numbers), then any space spanned by a subsequence is isomorphic to its square and hyperplanes. Furthermore, there is a subsequence of the basis such that all of its subsequences span isomorphic spaces.*

For example, as the usual basis of Tsirelson's space does not have a subsequence all of whose subsequences span isomorphic spaces, this shows that there is no isomorphic classification of the subspaces of Tsirelson's space by real numbers.

This result can be coupled with Gowers' dichotomy [6] proving:

Theorem 3 *Let X be a separable Banach space. Either E_0 Borel reduces to isomorphism between its subspaces or X contains a reflexive subspace with an unconditional basis all of whose subsequences span isomorphic spaces.*

For the above we will need some Ramsey type results for product spaces and some constructions for reducing E_0 . These results seem to have an independent interest apart from their applications to Banach space theory in that they classify minimal counter examples to Ramsey properties in product spaces. Let us just state one of these:

Theorem 4 *Let E be an analytic equivalence relation on $[\mathbb{N}]^{\mathbb{N}}$ invariant under finite changes. Either E_0 Borel reduces to E or E admits a homogeneous set.*

1.2 Schauder bases.

Let X be some separable Banach space and (e_i) a non zero sequence in X . We say that (e_i) is a *basis* for X if any vector x in X can be uniquely written as a norm convergent series $x = \sum a_i e_i$. In that case, the biorthogonal functionals $e_k^*(\sum a_i e_i) := a_k$ and the projections $P_n(\sum a_i e_i) := \sum_{i=0}^n a_i e_i$ are in fact continuous and moreover their norms are uniformly bounded.

If (e_i) is some non zero sequence that is a basis for its closed linear span, written $[e_i]$, we say that it is a *basic sequence* in X . The property of (e_i) being a basic sequence can also equivalently be stated as the existence of a constant $K \geq 1$ such that for any $n \leq m$ and $a_0, a_1, \dots, a_m \in \mathbb{R}$:

$$\left\| \sum_{i=0}^n a_i e_i \right\| \leq K \left\| \sum_{i=0}^m a_i e_i \right\|$$

Suppose furthermore that for any $x = \sum a_i e_i$ the series actually converges unconditionally, i.e., for any permutation σ of $\mathbb{N}$ the series $\sum a_{\sigma(i)} e_{\sigma(i)}$ converges to x . Then the basic sequence is said to be *unconditional*.

Again, being an unconditional basis for some closed subspace (which will be denoted by ‘unconditional basic sequence’) is equivalent to there being a constant $K \geq 1$, such that for all n , $A \subset \{0, \dots, n\}$ and $a_0, \dots, a_n \in \mathbb{R}$

$$\left\| \sum_{i \in A} a_i e_i \right\| \leq K \left\| \sum_{i=0}^n a_i e_i \right\|$$

We will in general only work with *normalised* basic sequences, i.e., $\|e_i\| \equiv 1$, which always can be obtained by taking $e'_i := \frac{e_i}{\|e_i\|}$.

Given some vector $x \in \text{span}(e_i)$ let its *support*, $\text{supp}(x)$, be the set of indices i with $e_i^*(x) \neq 0$. For $k \in \mathbb{N}$ and $x, y \in \text{span}(e_i)$ we write $k < x$ if

$k < \min \text{supp}(x)$ and $x < y$ if $\max \text{supp}(x) < \min \text{supp}(y)$. A *block basis*, (x_i) , over a basis (e_i) is a finite or infinite sequence of vectors in $\text{span}(e_i)$ with $x_0 < x_1 < x_2 < \dots$. This sequence will also be basic and in fact unconditional in case (e_i) is so.

Two basic sequences (e_i) and (t_i) are called *equivalent*, in symbols $(e_i) \approx (t_i)$, provided a series $\sum a_i e_i$ converges if and only if $\sum a_i t_i$ converges. This can also be stated as saying that $T: e_i \mapsto t_i$ extends to an invertible linear operator between $[e_i]$ and $[t_i]$. The quantity $\|T\| \cdot \|T^{-1}\|$ is then the constant of equivalence between the two bases.

A basis that is equivalent to all of its subsequences is said to be *sub-symmetric*. A simple diagonalisation argument then shows that it must be uniformly equivalent to all of its subsequences.

Two basic sequences (e_i) and (t_i) are said to be *permutatively equivalent* if there is some permutation σ of $\mathbb{N}$ such that (e_i) and $(t_{\sigma(i)})$ are equivalent.

2 Incomparable and minimal subspaces.

Two Banach spaces X and Y are called *incomparable* in case neither of them embed isomorphically into the other. X is said to be *minimal* if it embeds into all of its infinitely dimensional subspaces and X itself is infinite dimensional.

Our proof of the first theorem will proceed by a reduction to an analysis of Borel partial orders due to L. Harrington, D. Marker and S. Shelah (see [9]). Instrumental in our reduction will be the determinacy result of Gowers on certain games in Banach spaces (see [6]), which will guarantee that some choices can be done uniformly, a fact that is needed for definability purposes. Moreover, we will use some ideas of J. Lopez-Abad on coding reals with inevitable subsets of the unit sphere of a Banach space (see [14]).

We mention that it was shown by a simpler argument in [5] by V. Ferenczi and the author that any Banach space either contains a minimal subspace or a continuum of non isomorphic subspaces.

For facility of notation we write $X \sqsubseteq Y$ if X embeds isomorphically into Y and will always suppose the spaces we are working with to be separable infinite dimensional. Then $\sqsubseteq$ restricted to the standard Borel space of subspaces of some separable Banach space becomes an analytic quasi-order, i.e., transitive and reflexive. So the result above amounts to saying that either $\sqsubseteq$ has a minimal element or a perfect antichain.

Suppose (e_i) is a normalised basic sequence with norm denoted by $\|\cdot\|$.

We call a normalised block vector x with finite support rational if it is a scalar multiple of a finite linear combination of (e_i) with rational coordinates. Notice that there are only countably many rational (finite) block vectors, which we can gather in a set Q and give it the discrete topology. Let $bb_{\mathbb{Q}}(e_i)$ be the set of block bases of (e_i) consisting of rational normalised block vectors, which is easily seen to be a closed subspace of $Q^{\mathbb{N}}$, which is itself a Polish space. Moreover the canonical function sending $X \in bb_{\mathbb{Q}}(e_i)$ to its closed span in $\mathfrak{B}$ is Borel, so the relations of isomorphism, etc., become analytic on $bb_{\mathbb{Q}}(e_i)$.

We recall the following classical facts: Any infinite dimensional Banach space contains an infinite normalised basic sequence (e_i) . Moreover, if Y is any subspace of $[e_i]$, then it contains an isomorphic perturbation of a block basic sequence of (e_i) . Again any block basic sequence is equivalent to some member of $bb_{\mathbb{Q}}(e_i)$. So this explains why we can concentrate on $bb_{\mathbb{Q}}(e_i)$ if we are only looking for minimal subspaces.

For $X, Y \in bb_{\mathbb{Q}}(e_i)$, let $X \leq Y$ if X is a blocking of Y , i.e., if any element of X is a linear combination over Y . Note that this does not imply that they are rational block vectors over Y , but only over (e_i) . Moreover, if $Y = (y_i), X = (x_i) \in bb_{\mathbb{Q}}(e_i)$, put $Y \leq^* X$ if for some k , $(y_i)_{i \geq k} \leq X$. Also, for $\Delta = (\delta_i)$ an infinite sequence of strictly positive reals write $d(X, Y) < \Delta$ if $\forall i \ \|x_i - y_i\| < \delta_i$.

Put $X \approx Y$ if the bases are equivalent and $X \cong Y$ if they span isomorphic spaces. Then a classical perturbation argument shows that there is some Δ depending only on the constant of the basis, such that for any $X, Y \in bb_{\mathbb{Q}}(e_i)$ if $d(X, Y) < \Delta$, then $X \approx Y$ and in particular $X \cong Y$. Put also $X = (x_i) \simeq Y = (y_i)$ if $\exists k \ \forall i \geq k \ x_i = y_i$. Then evidently $X \simeq Y$ implies $X \approx Y$.

For a subset $A \subset bb_{\mathbb{Q}}(e_i)$ let $A^* = \{Y \in bb_{\mathbb{Q}}(e_i) \mid \exists X \in A \ X \simeq Y\}$ and $A_{\Delta} = \{Y \in bb_{\mathbb{Q}}(e_i) \mid \exists X \in A \ d(X, Y) < \Delta\}$. Notice that if A is analytic so are both A^* and A_{Δ} . Again $[Y] = \{X \in bb_{\mathbb{Q}}(e_i) \mid X \leq Y\}$. Such an A is said to be large in $[Y]$ if for any $X \in [Y]$ we have $[X] \cap A \neq \emptyset$.

For (e_i) a given normalised basis, $A \subset bb_{\mathbb{Q}}(e_i)$ and $X \in bb_{\mathbb{Q}}(e_i)$, the Gowers game $\mathcal{D}_X^A$ is defined as follows: Player I plays in the k 'th move of the game a rational normalised block vector y_k of (e_i) such that $y_{k-1} < y_k$ and y_k is a block on X . Player II responds by either doing nothing or playing a rational normalised block vector x such that $x \in [y_{l+1}, \dots, y_k]$ where l was the last move where II played a vector. So player II wins the game if in the end she has produced an infinite rational block basis $X = (x_i) \in A$. This is an equivalent formulation due to J. Bagaria and J. Lopez-Abad (see [1]) of

Gowers' original game.

Gowers [6] proved that if $A \subset bb_{\mathbb{Q}}(e_i)$ is analytic, large in $[Y]$ and Δ is given, then for some $X \in [Y]$ II has a winning strategy in the game $\mathfrak{D}_X^{A\Delta}$.

We mention also a result of Odell and Schlumprecht [15] obtained from their solution to the distortion problem: If E is an infinite dimensional Banach space not containing c_0 , there are an infinite dimensional subspace F and $A, B \subset \mathcal{S}_F$ of positive distance such that any infinite dimensional subspace of F intersects both A and B .

The following was shown in [5]:

Lemma 5 (MA) *Let $A \subset bb_{\mathbb{Q}}(e_i)$ be linearly ordered under $\le^*$ of cardinality strictly less than the continuum. Then there is some $X \in bb_{\mathbb{Q}}(e_i)$ such that $X \le^* Y$ for all $Y \in A$.*

From this lemma one gets the following:

Lemma 6 (MA + $\neg CH$) *Suppose $W \subset bb_{\mathbb{Q}}(e_i)$ is a Σ_2^1 set, large in some $[Y]$ and $\Delta > 0$. Then II has a winning strategy in $\mathfrak{D}_X^{W\Delta}$ for some $X \in [Y]$.*

Proof : Let $W = \bigcup_{\omega_1} V_\xi$ be a decomposition of W as an increasing union of $\aleph_1$ Borel sets. We claim that some V_ξ^* is large in $[Z]$ for some $Z \in [Y]$, which by Gowers' theorem will be enough to prove the lemma. So suppose not and find $Y_0 \in [Y]$ such that $[Y_0] \cap V_0^* = \emptyset$. Repeating the same process and diagonalising at limits, we find $Y_\xi \in [Y]$ for $\xi < \omega_1$ such that $[Y_\xi] \cap V_\xi^* = \emptyset$ and $Y_\xi \le^* Y_\eta$ for $\eta < \xi$. By the above lemma there is some $Y_\infty = (y_i) \in [Y]$ with $Y_\infty \le^* Y_\xi$ for all $\xi < \omega_1$.

We claim that $[(y_{2i})] \cap W = \emptyset$. Otherwise, for $Z = (z_i) \in [(y_{2i})] \cap W$ find $\xi < \omega_1$ such that $Z \in V_\xi$. Now as $(y_i) \le^* Y_\xi$ there is some k with $(y_i)_{i \geq k} \leq Y_\xi$, but then

$$(y_{2i}) \simeq (y_k, y_{k+1}, y_{k+2}, \dots, y_{2k-1}, y_{2k}, y_{2(k+1)}, y_{2(k+2)}, \dots) \leq Y_\xi$$

One now easily sees that there is some (x_i) with

$$(z_i) \simeq (x_i) \leq (y_k, y_{k+1}, y_{k+2}, \dots, y_{2k-1}, y_{2k}, y_{2(k+1)}, y_{2(k+2)}, \dots)$$

whereby $(x_i) \in V_\xi^*$ contradicting $V_\xi^* \cap [Y_\xi] = \emptyset$.

Therefore $[(y_{2i})] \cap W = \emptyset$, again contradicting the largeness of W . $\square$

Lemma 7 ($MA + \neg CH$) *Suppose that (e_i) is a basic sequence such that $[e_i]$ does not contain a minimal subspace. Then for any $Y \in bb_{\mathbb{Q}}(e_i)$ there are a $Z \in [Y]$ and a Borel function $g: [Z] \rightarrow [Z]$, with $g(X) \leq X$ and $X \not\sqsubseteq g(X)$ for all $X \in [Z]$.*

Proof: As c_0 is minimal, $[e_i]$ does not contain c_0 . Therefore, by the solution to the distortion problem by Odell and Schlumprecht, we can by replacing (e_i) by a block suppose that we have two positively separated sets F_0, F_1 of the unit sphere, such that for any $X \in bb_{\mathbb{Q}}(e_i)$ there are rational normalised blocks x, y on X with $x \in F_0$ and $y \in F_1$. We call such sets inevitable.

Let $D = \{X = (x_i) \in bb_{\mathbb{Q}}(e_i) \mid \forall i \ x_i \in F_0 \cup F_1\}$ and for $X \in D$ let $\alpha(X) \in 2^{\mathbb{N}}$ be defined by $\alpha(X)(i) = 0 \leftrightarrow x_i \in F_0$. Then D is easily seen to be a closed subset of $bb_{\mathbb{Q}}(e_i)$ and $\alpha: D \rightarrow 2^{\mathbb{N}}$ to be continuous. Furthermore by the inevitability of F_0 and F_1 we have that D is large in every $[Y]$.

Let $\mathbb{Q}_*^{<\mathbb{N}}$ be the set of finite non identically zero sequences of rational numbers given the discrete topology. Then $(\mathbb{Q}_*^{<\mathbb{N}})^{\mathbb{N}}$ is Polish. Define for any $Y \in bb_{\mathbb{Q}}(e_i)$ and $(\underline{\lambda}_i) \in (\mathbb{Q}_*^{<\mathbb{N}})^{\mathbb{N}}$ the block basis $(\underline{\lambda}_i) \cdot Y$ of Y in the obvious way, by taking the linear combinations given by $(\underline{\lambda}_i)$.

Fix also some perfect set P of almost disjoint subsets of $\mathbb{N}$ seen as a subset of $2^{\mathbb{N}}$ and let $\beta: P \leftrightarrow (\mathbb{Q}_*^{<\mathbb{N}})^{\mathbb{N}}$ be a Borel isomorphism.

Again $E = \{X \in D \mid \alpha(X) \in P\}$ is large and closed in $bb_{\mathbb{Q}}(e_i)$.

Then the set

$$W = \{X = (x_i) \in bb_{\mathbb{Q}}(e_i) \mid (x_{2i}) \in E \wedge (x_{2i+1}) \not\sqsubseteq \beta \circ \alpha((x_{2i})) \cdot (x_{2i+1})\}$$

is coanalytic. We claim moreover that it is large in $bb_{\mathbb{Q}}(e_i)$.

To see this, let $Y \in bb_{\mathbb{Q}}(e_i)$ be given and take by inevitability of F_0 and F_1 some $(z_i) \in [Y]$ with $z_{3i} \in F_0$ and $z_{3i+1} \in F_1$. As $[z_{3i+2}]$ is not minimal there is some $X \leq (z_{3i+2})$ such that $(z_{3i+2}) \not\sqsubseteq X$. Take some $(\underline{\lambda}_i) \in (\mathbb{Q}_*^{<\mathbb{N}})^{\mathbb{N}}$ such that $(\underline{\lambda}_i) \cdot (z_{3i+2}) \approx X$ and $(z_{3i+2}) \not\sqsubseteq (\underline{\lambda}_i) \cdot (z_{3i+2})$. We can now define some (v_i) such that either $v_{2i} = z_{3i}$ or $v_{2i} = z_{3i+1}$, $\beta \circ \alpha((v_{2i})) = (\underline{\lambda}_i)$ and $v_{2i+1} = z_{3i+2}$. This ensures that $(v_i) \in W$. So as $(v_i) \leq (z_i)$ it is in $[Y]$ and W is indeed large.

Take now some $\Delta = (\delta_i)$ depending on the basic constant as above with $\delta_i < \frac{1}{2}d(F_0, F_1)$. By the preceding lemma we can find a $Y \in bb_{\mathbb{Q}}(e_i)$ such that II has a winning strategy σ in the game $\mathfrak{D}_Y^{W^* \Delta}$.

Suppose that (x_i) has been played by II according to the strategy σ as a response to Z played by I. As σ is winning, $X \in W_{\Delta}^*$. Define $\gamma(X) \in 2^{\mathbb{N}}$

by $\gamma(X)(i) = 0$ if $d(x_{2i}, F_0) < \delta_{2i}$ and $\gamma(X)(i) = 1$ otherwise. Then γ is Borel from W_Δ^* to $2^\mathbb{N}$, and furthermore there is a unique $\gamma^*(X) \in P$ such that $\exists k \forall i \geq k \gamma(X)(i) = \gamma^*(X)(i)$. This is because P was chosen to consist of almost disjoint subsets of $\mathbb{N}$. Again $X \mapsto \gamma^*(X)$ is Borel.

Take some $U = (u_i) \in W$ such that $\forall^\infty n \|u_n - x_n\| < \delta_n$. Then $\alpha(U) = \gamma^*(X)$, $(u_{2i+1}) \approx (x_{2i+1})$ and $(u_{2i+1}) \not\sqsubseteq \beta \circ \alpha(U) \cdot (u_{2i+1})$. So due to the equivalence invariance of the basis by Δ perturbations we have $(x_{2i+1}) \not\sqsubseteq \beta \circ \gamma^*(X) \cdot (x_{2i+1})$.

Let $V \in [X]$ be the normalisation of $\beta \circ \gamma^*(X) \cdot (x_{2i+1})$. The function $g: Z \mapsto V$ is Borel and obviously

$$V \approx \beta \circ \gamma^*(X) \cdot (x_{2i+1}) \leq (x_{2i+1}) \leq Z$$

and as

$$(x_{2i+1}) \not\sqsubseteq \beta \circ \gamma^*(X) \cdot (x_{2i+1})$$

also $Z \not\sqsubseteq V$. □

A Banach space is called quasi-minimal if any two subspaces have further isomorphic subspaces. The following is a standard observation.

Lemma 8 *Suppose $[e_i]$ is quasi-minimal. Then $\sqsubseteq$ is downwards σ -directed on $bb_\mathbb{Q}(e_i)$, i.e., any countable family has a common minorant.*

Proof: Suppose that $Y_i \in bb_\mathbb{Q}(e_i)$ are given, then define inductively $Z_i \in [Y_0]$ such that $Z_i \sqsubseteq Y_i$ and $Z_{i+1} \leq Z_i$. Take some $Z = (z_i) \leq^* Z_n$ for all n and notice as in the proof of lemma 6 that $(z_{2i}) \sqsubseteq Z_n$ for all n . □

Lemma 9 *If R is a downwards σ -directed Borel quasi-order on a standard Borel space X . Then either R has a perfect antichain or a minimal element.*

Proof: This is a simple consequence of the results of L. Harrington, D. Marker and S. Shelah [9], as we will see. Suppose that R did not have a perfect antichain, then by their results there is a countable partition $X = \bigcup X_n$ into Borel sets, so that R is total on each piece, i.e., R can be written as a countable union of R -chains.

Applying another of their results this implies that for some countable ordinal α there are Borel functions $f_n: X_n \rightarrow 2^\alpha$, such that for any $x, y \in X_n$:

$$yRx \iff x \leq_{lex} y$$

Where $\leq_{lex}$ is the usual lexicographical ordering. In their terminology, R is linearisable on each X_n .

One can easily check that any subset of 2^α has a countable subset cofinal with respect to $\leq_{lex}$, so pulling it back by f_n it becomes coinitial in $R \upharpoonright_{X_n}$. Putting all these sets together one gets a countable subset of X coinitial with respect to R . So by downwards σ -directedness there is therefore a minimal element in X . $\square$

After this series of lemmas we can now prove the theorem:

Theorem 10 *Let X be an infinite dimensional Banach space. Then X contains either a minimal subspace or a continuum of pairwise incomparable subspaces.*

Proof : By Gowers' quadrichotomy X contains either a quasi-minimal subspace or a subspace with a basis such that any two disjointly supported subspaces are totally incomparable (see Gowers [6] theorem 7.2 and the fact that H.I. spaces are quasi-minimal). In the latter case any perfect set of almost disjoint subsets of $\mathbb{N}$ will give rise to subsequences of the basis spanning totally incomparable spaces, which would prove the theorem. So we can suppose that $X = [e_i]$ is quasi-minimal for some basis (e_i) . If X does not contain a minimal subspace, we can choose $Z \in bb_{\mathbb{Q}}(e_i)$ and the Borel function as above (under $\mathbf{MA} + \neg\mathbf{CH}$ of course). So define the following property on subsets A, B of $[Z]^2$:

$$\Phi(A, B) \Leftrightarrow$$

$$\forall Y, V, W \in [Z] \left[(Y, V) \notin A \vee (V, W) \notin A \vee (Y, W) \notin B \right] \wedge \forall Y \in [Z] (Y, g(Y)) \notin A$$

We see that Φ is Π_1^1 on Σ_1^1 , hereditary and continuous upwards in the second variable. Furthermore, $\Phi(\sqsubseteq, \not\sqsubseteq)$, so by the second reflection theorem (see Kechris [11] theorem (35.16)) there is some Borel set R containing $\sqsubseteq$ such that $\Phi(R, \mathbb{C}R)$. But then R is a Borel quasi-order, downwards σ -directed, as it contains $\sqsubseteq$, and without a minimal element, as witnessed by g . So R has a perfect anti-chain by the previous lemma, which then is an antichain for $\sqsubseteq$ too.

The statement is therefore proved under the additional hypothesis of Martin's axiom and the negation of the continuum hypothesis. We will see that this is in fact sufficient to prove the theorem. By standard metamathematical facts and Shoenfield's absoluteness theorem it is enough to show that the statement we wish to prove is Σ_2^1 .

It was proved by Ferenczi and the author in [5] that the property of having a block minimal subspace was Σ_2^1 . For using Gowers' determinacy result and codings as above, one can continuously find an isomorphism between the space and a certain subspace to testify the minimality. This proof can trivially be modified to show that the property of having a minimal (i.e., not necessarily block minimal) subspace is also Σ_2^1 . For now we only have to choose not a code for a subspace and an isomorphism, but a code for a subspace and an embedding. For the convenience of the reader, we have included the proof of this in an appendix.

On the other hand, the property of having a perfect antichain is obviously Σ_2^1 by just counting quantifiers. So these remarks finish the proof. $\square$

3 Ramsey type results.

We will show two Ramsey type results and afterwards some applications to Banach space theory.

It is well known that there are no nice Ramsey properties for the product space $[\mathbb{N}]^{\mathbb{N}} \times [\mathbb{N}]^{\mathbb{N}}$ in contradistinction to the simple Ramsey space $[\mathbb{N}]^{\mathbb{N}}$. That is, there are even quite simple relations not admitting a square $[A]^{\mathbb{N}} \times [B]^{\mathbb{N}}$ that is either included in or disjoint from the relation. An example of this is the oscillation relation $\mathcal{O}$ defined by

$$(a_n)\mathcal{O}(b_n) \iff \exists N \forall n \left[\#(k \mid a_n < b_k < a_{n+1}) \leq N \wedge \#(k \mid b_n < a_k < b_{n+1}) \leq N \right]$$

Where $[\mathbb{N}]^{\mathbb{N}}$ is seen as the space of strictly increasing sequences of integers (a_i) .

The situation is very different if one replaces one of the factor spaces by other Ramsey spaces and there are now very deep positive theorems on so called polarised partition relations.

We are interested in the case when the relation on the product is in fact a definable equivalence relation. Here the right question seems to be when there is a cube $[A]^{\mathbb{N}}$ contained in one class. Now if one lets two subsets of $\mathbb{N}$ be equivalent iff they have the same minimal element, then the relation has exactly $\aleph_0$ classes and does not admit a homogeneous set.

On the other hand if the relation is invariant under finite changes, such as E_0 , then there are bigger chances that it should have a homogeneous set.

We will show that in the case of analytic equivalence relations, E_0 is in fact the minimal counterexample to the Ramsey property, in the sense that, if an analytic equivalence relation is invariant under finite changes and does not admit a homogeneous set, then it Borel reduces E_0 . In the same vein it is shown that if an analytic equivalence relation does not admit a cube on which it has only countably many classes, then it has at least a perfect set of classes. We notice that both of these results are relatively direct consequences of the Silver and Glimm-Effros dichotomies in the case of the equivalence relation being Borel. But our results are motivated by applications to isomorphism of separable Banach spaces, which is true analytic, and the dichotomies are known not to hold in this generality.

The following result was also found independently by S. Todorcevic, albeit with a somewhat different proof:

Theorem 11 *Let E be an analytic equivalence relation on $[\mathbb{N}]^{\mathbb{N}}$. Then either E has a continuum of classes or there is some $A \in [\mathbb{N}]^{\mathbb{N}}$ such that E only has a countable number of classes on $[A]^{\mathbb{N}}$.*

Moreover, A 's E_0 class will be a complete section for E on $[A]^{\mathbb{N}}$.

Proof : We will prove the theorem under $\mathbf{MA} + \neg\mathbf{CH}$. By Burgess' theorem (Exercise (35.21) in [12]) we can suppose that E has at most $\aleph_1$ classes $(C_\xi)_{\omega_1}$. Define $P_\xi(A) \leftrightarrow [A]_{E_0} \cap C_\xi \neq \emptyset$ and notice that this is an analytic E_0 -invariant property. We can by simple diagonalisation find $(A_\xi)_{\omega_1}$, $A_\xi \subset^* A_\eta$ for $\eta < \xi < \omega_1$ such that $\forall \xi < \omega_1$ either $[A_\xi]^{\mathbb{N}} \subset P_\xi$ or $[A_\xi]^{\mathbb{N}} \subset \mathbb{C}P_\xi$. And by $\mathbf{MA} + \neg\mathbf{CH}$ there is an $A \subset^* A_\xi$, $\forall \xi < \omega_1$.

Notice now that by E_0 -invariance of P_ξ , if $B \subset^* A$ and $[A]^{\mathbb{N}} \subset P_\xi$ or $[A]^{\mathbb{N}} \subset \mathbb{C}P_\xi$ then also $[B]^{\mathbb{N}} \subset P_\xi$, respectively $[B]^{\mathbb{N}} \subset \mathbb{C}P_\xi$. So therefore $\forall \xi < \omega_1$ $[A]^{\mathbb{N}} \subset P_\xi$ or $[A]^{\mathbb{N}} \subset \mathbb{C}P_\xi$.

Suppose now that $B \in [A]^{\mathbb{N}}$, $B \in C_\xi$ then $P_\xi(B)$ and therefore $[A]^{\mathbb{N}} \subset P_\xi$ and $P_\xi(A)$, i.e., $\exists A' E_0 A \quad A' E B$. This means that $[A]_{E_0}$ is a complete section for E on $[A]^{\mathbb{N}}$.

Let us now see that the statement of the theorem is absolute. Saying that E has a continuum of classes is equivalent to saying that there is a compact perfect set $K \subset [\mathbb{N}]^{\mathbb{N}}$ consisting of pairwise E -inequivalent points:

$$\exists K \subset [\mathbb{N}]^{\mathbb{N}} \text{ compact, perfect } \forall x, y \in K (x = y \vee x E y)$$

This is obviously a Σ_2^1 statement.

For the other case, notice that as $[A]_{E_0}$ is a complete section for E on $[A]^\mathbb{N}$ there is by the Jankov-von Neumann selection theorem a C -measurable selector $f: [A]^\mathbb{N} \rightarrow \mathbb{N}^\mathbb{N}$ and a Borel set $\mathcal{B} \subset [\mathbb{N}]^\mathbb{N} \times [\mathbb{N}]^\mathbb{N} \times \mathbb{N}^\mathbb{N}$ with $E = \pi_{[\mathbb{N}]^\mathbb{N} \times [\mathbb{N}]^\mathbb{N}} \mathcal{B}$, such that for $D \in [A]^\mathbb{N}$ there is an $A'E_0A$ with $\mathcal{B}(D, A', f(D))$. That is, we can choose a witness to D being E equivalent to some $A'E_0A$ in a C -measurable way. But any C -measurable function can, using Ellentuck's theorem, be rendered continuous on a cube, i.e., there is some $B \in [A]^\mathbb{N}$ such that f 's restriction to $[B]^\mathbb{N}$ is continuous. So by the proof above the E -classes on $[A]^\mathbb{N}$ are the same as the E -classes on $[B]^\mathbb{N}$ and the other possibility can be written as:

$$\exists B \exists f: [B]^\mathbb{N} \rightarrow \mathbb{N}^\mathbb{N} \text{ continuous } \forall D \in [B]^\mathbb{N} \exists B' E_0 B \mathcal{B}(D, B', f(D))$$

This statement is Σ_2^1 as the quantifier $\exists B' E_0 B$ is over a countable set, so by Shoenfield absoluteness and standard metamathematical facts it is enough to prove the result under $\mathbf{MA} + \neg \mathbf{CH}$. $\square$

Our next results render explicit the connection with the Borel reducibility ordering.

Definition 12 For $A, B \subset \mathbb{N}$ set AE'_0B iff $\exists n |A \cap n| = |B \cap n| \wedge A \setminus n = B \setminus n$

It is easy to see that the equivalence class of any infinite-cofinite subset of $\mathbb{N}$ is dense in $[\mathbb{N}]^\mathbb{N}$ and in fact the equivalence relation is generically ergodic. Moreover, E'_0 is just a refinement of E_0 .

Lemma 13 E'_0 is generically ergodic (i.e., any invariant set with the Baire property is either meagre or comeagre) and all classes $[A]_{E'_0}$, for A infinite-cofinite, are dense.

Proof : Since $[\mathbb{N}]^\mathbb{N}$ is cocountable in $2^\mathbb{N}$ we can restrict our attention to it. Suppose that some invariant set $\mathcal{A}$ is non meagre, then there is some $a \subset [0, n]$ such that $\mathcal{A}$ is comeagre in $D_{a,n} = \{A \in [\mathbb{N}]^\mathbb{N} \mid A \cap [0, n] = a\}$. So for any $D_{b,m}$ there are $c, d \in [0, k]$; $\max(n, m) < k$ such that $a \subset c$, $b \subset d$, $|c| = |d|$. Now for any $A \in [\{k+1, k+2, \dots\}]^\mathbb{N}$ we have $\phi(c \cup A) := (d \cup A)E'_0(c \cup A)$ and ϕ is a homeomorphism of $D_{c,k} \subset D_{a,n}$ with $D_{d,k}$. But that means that the image of $\mathcal{A}$ is comeagre in $D_{d,k} \subset D_{b,m}$ and is included in the saturation of $\mathcal{A}$, which is $\mathcal{A}$. So $\mathcal{A}$ is comeagre in the space.

If A is infinite-cofinite, then for any $D_{a,n}$ there are $b, c \subset [0, k]$; $b \supset a$, $n < k$, $A \cap [0, k] = c$, $b \cap [0, n] = a$ and $|b| = |c|$. So $A = (c \cup A/k)E'_0(b \cup A/k) \in D_{b,k} \subset D_{a,n}$. And its class is dense. $\square$

Proposition 14 *Let E be a meagre equivalence relation on $2^{\mathbb{N}}$ containing E'_0 . Then $E_0 \leq_B E$.*

Proof : Let (D_n) be a decreasing sequence of dense open sets, such that $E \cap \bigcap_n D_n = \emptyset$.

We will inductively construct sequences $b_0^n, b_1^n \in 2^{<\mathbb{N}}$ for $n \in \mathbb{N}$ such that for all n , $|b_0^n| = |b_1^n|$, $\overline{b_0^n} = \overline{b_1^n} := \#\{k \mid b_1^n(k) = 1\}$. And if $a_s := b_{s(0)}^0 \wedge \dots \wedge b_{s(|s|-1)}^{|s|-1}$ for all $s \in 2^{<\mathbb{N}}$, then for any $s, t \in 2^n$, $N_{a_s \wedge 0} \times N_{a_t \wedge 1} \subset D_{n+1}$.

Suppose that this can be done. Then define $\alpha \mapsto \bigcup_n a_{\alpha \upharpoonright n} = a_\alpha$. This is clearly continuous. If now $\neg \alpha E_0 \beta$, then for infinitely many n , $\alpha(n) \neq \beta(n)$. So for these n $(a_\alpha, a_\beta) \in N_{a_{\alpha \upharpoonright n+1}} \times N_{a_{\beta \upharpoonright n+1}} \subset D_{n+1}$, which implies that $(a_\alpha, a_\beta) \in \bigcap_k D_k \subset \mathbb{C}E$.

Conversely, if $\alpha E_0 \beta$, then for some N , we have $\forall n \geq N \alpha(n) = \beta(n)$. But then easily $a_\alpha = a_{\alpha \upharpoonright N} \wedge b_{\alpha(N)}^N \wedge b_{\alpha(N+1)}^{N+1} \dots$ and $a_\beta = a_{\beta \upharpoonright N} \wedge b_{\alpha(N)}^N \wedge b_{\alpha(N+1)}^{N+1} \dots$, so by the construction, $a_\alpha E'_0 \beta$.

Now for the construction: Suppose that b_0^n, b_1^n have been chosen for $\forall m < n$, enumerate $2^n \times 2^n$ by $(s_0, t_0), \dots, (s_k, t_k)$ and take $c_0^0, c_1^0 \in 2^{<\mathbb{N}}$ such that $N_{a_{s_0} \wedge c_0^0} \times N_{a_{t_0} \wedge c_1^0} \subset D_n$. This can be done as D_n is dense and open in the product.

Prolong c_0^0, c_1^0 to c_0^1, c_1^1 respectively in such a way that $N_{a_{s_1} \wedge c_0^1} \times N_{a_{t_1} \wedge c_1^1} \subset D_n$.

Again, prolong c_0^1, c_1^1 to c_0^2, c_1^2 respectively in such a way that $N_{a_{s_2} \wedge c_0^2} \times N_{a_{t_2} \wedge c_1^2} \subset D_n$, etc.

Finally, prolong c_0^k, c_1^k to b_0^n, b_1^n respectively, such that $|b_0^n| = |b_1^n|$, $\overline{b_0^n} = \overline{b_1^n}$. This finishes the construction. $\square$

For the following, we recall that $\forall^* x R(x)$ means that the set $\{x \mid R(x)\}$ is comeagre, where x varies over some Polish space.

Theorem 15 *Let E be an analytic equivalence relation on $[\mathbb{N}]^{\mathbb{N}}$ such that $E'_0 \subset E$, i.e., E is E'_0 -invariant. Then either $E_0 \leq_c E$ or there is some $A \in [\mathbb{N}]^{\mathbb{N}}$ such that E only has one class on $[A]^{\mathbb{N}}$.*

Proof : By corollary 3.5 of [10], if $E_0 \not\leq_c E$, then E will be a decreasing intersection of $\aleph_1$ smooth equivalence relations:

$$E = \bigcap_{\omega_1} E_\xi, \quad E_\xi \subset E_\eta, \quad \eta < \xi < \omega_1$$

Let $f_\xi: [\mathbb{N}]^\mathbb{N} \rightarrow \mathbb{R}$ be a Borel reduction of E_ξ to identity on $\mathbb{R}$. Then for any $A \in [\mathbb{N}]^\mathbb{N}$, there is a $B \in [A]^\mathbb{N}$ such that $f \upharpoonright [B]^\mathbb{N}$ is continuous. But since there is a dense E_ξ -class the function has to be constant, that is, there is only one class.

We construct inductively a $\subset^*$ -decreasing sequence $(A_\xi)_{\omega_1}$ of infinite subsets of $\mathbb{N}$, with each A_ξ being homogeneous for E_ξ . Under **MA** + $\neg\mathbf{CH}$ such a sequence can be diagonalised to produce an infinite $A_\infty \subsetneq^* A_\xi$, $\forall \xi < \omega_1$. Now as $A_\infty \subsetneq^* A_\xi$ it is easily seen that A_∞ is E'_0 -equivalent with some subset of A_ξ and therefore also E_ξ -equivalent with A_ξ itself. Furthermore, the same holds for any infinite subset of A_∞ , so A_∞ is homogeneous for all of the E_ξ and therefore for E too.

As before one sees that the property of having a homogeneous set is Σ_2^1 , so we need only check that continuously reducing E_0 is Σ_2^1 . But this can be written as:

$$\begin{aligned} \exists f: [\mathbb{N}]^\mathbb{N} \rightarrow [\mathbb{N}]^\mathbb{N} \text{ continuous } & \left[\forall^* \alpha \in [\mathbb{N}]^\mathbb{N} \forall \beta E_0 \alpha \forall \gamma E_0 \alpha f(\beta) E f(\gamma) \right. \\ & \left. \wedge \forall \alpha, \beta \in [\mathbb{N}]^\mathbb{N} \{ \alpha E_0 \beta \vee \neg f(\alpha) E f(\beta) \} \right] \end{aligned}$$

So as the quantifier $\forall \beta E_0 \alpha$ is over a countable set and that the category quantifier $\forall^*$ preserves analyticity (see Theorem (29.22) in [12]), the statement is Σ_2^1 . $\square$

4 Applications to Banach space theory.

Let (e_i) be some basic sequence in a Banach space X and define the following equivalence relation on $[\mathbb{N}]^\mathbb{N}$: $A \cong B \iff [e_i]_A \cong [e_i]_B$. Then $\cong$ is analytic and extends E'_0 . For suppose that $AE'_0 B$. Then $[e_i]_A$ and $[e_i]_B$ are spaces of the same finite codimension in $[e_i]_{A \cup B}$ and are therefore isomorphic. So, using the proposition, one sees that if $E_0 \not\leq_B \cong$, then $\cong$ must be non meagre and therefore by Kuratowski-Ulam have a non meagre class, which again by the lemma is comeagre.

To avoid trivialities, let us in the following suppose that all Banach spaces considered are separable, infinite dimensional.

Gowers showed the following amazing result about the structure of subspaces of a Banach space: if X is a Banach space, then it contains either an unconditional basic sequence or an H.I. subspace [6].

Here an H.I. (hereditarily indecomposable) space Y is one in which no two infinite dimensional subspaces form a direct sum. This property, which passes to subspaces, insures that Y cannot be isomorphic to any of its subspaces and cannot contain any unconditional basic sequence. Therefore in the classification of the subspaces of a Banach space one can always suppose to be dealing with an H.I. space or a space with an unconditional basis.

Proposition 16 *Let (e_i) be a basic sequence in a Banach space. Then either E_0 Borel reduces to isomorphism of spaces spanned by subsequences of the basis or there will be some infinite $A \subset \mathbb{N}$, such that for any infinite $B \subset A$: $[e_i]_A \cong [e_i]_B$.*

Proof : This follows from Theorem 15 as isomorphism is E'_0 -invariant. $\square$

Example 17 *Hereditarily Indecomposable spaces.*

Suppose that we are given a hereditarily indecomposable space X . Then as any Banach space contains a (conditional) basic sequence, we can suppose that we have a basis (e_i) . By the above proposition, if E_0 does not reduce, there would be a subsequence spanning a space isomorphic to some proper subspace in contradiction with the properties of H.I. spaces. So E_0 reduces to isomorphism of its subspaces. The same reasoning shows, using the first theorem, that it has a continuum of incomparable subspaces.

A recent result due to Ferenczi and Galego [3] says that E_0 Borel reduces to the isomorphism relation between subspaces of c_0 and ℓ_1 . So if E_0 does not reduce to isomorphism between the subspaces of an Banach space, then using Gowers' dichotomy we can find a subspace with an unconditional basis. Therefore by James' characterisation of reflexivity this basis must span a reflexive space. All in all this gives us the following:

Theorem 18 *Let X be an Banach space such that the isomorphism relation between its subspaces does not reduce E_0 . Then X contains a reflexive subspace with an unconditional basis, all of whose subsequences span isomorphic spaces.*

Let us notice that if a basis (e_i) has the property that no two disjointly supported block basic sequences are equivalent, then one can easily show that this basis has the Casazza property and moreover that it satisfies

$$(e_i)_A \approx (e_i)_B \iff [e_i]_A \cong [e_i]_B \iff AE'_0B$$

See the work of Gowers and Maurey, [7], for unconditional examples of such bases. So as E'_0 and E_0 are Borel bi-reducible, there are bases on which both equivalence and isomorphism between subsequences are exactly of complexity E_0 .

We will now see an extension of some results by Ferenczi and the author, [4], and Kalton, [11].

Theorem 19 *Let (e_i) be an unconditional basic sequence. Then either E_0 Borel reduces to isomorphism of spaces spanned by subsequences of the basis or any space spanned by a subsequence is isomorphic to its square and its hyperplanes. And there is some infinite $A \subset \mathbb{N}$ such that for any infinite $B \subset A$, $[e_i]_A \cong [e_i]_B$.*

Proof : As before we can suppose we have some comeagre class $\mathcal{A} \subset [\mathbb{N}]^{\mathbb{N}}$. But then $\mathcal{A}$ is also comeagre in $2^{\mathbb{N}}$ and there is therefore a partition A_0, A_1 of $\mathbb{N}$ and subsets $B_0 \subset A_0, B_1 \subset A_1$ such that for any $C \subset \mathbb{N}$, if $C \cap A_0 = B_0$ or $C \cap A_1 = B_1$, then $C \in \mathcal{A}$. In particular, $B_0, B_1, B_0 \cup B_1 \in \mathcal{A}$. Moreover, as the complement operation is a homeomorphism of $2^{\mathbb{N}}$ with itself, there is some C such that $C, \mathbb{C}C \in \mathcal{A}$. So identifying subsets of $\mathbb{N}$ with the Banach spaces they generate and using the fact that the basis is unconditional, and therefore that disjoint subsets form direct sums, we can calculate:

$$\mathbb{N} = C \cup \mathbb{C}C \cong C \oplus \mathbb{C}C \cong B_0 \oplus B_0 \cong B_0 \oplus B_1 \cong B_0 \cup B_1 \cong B_0$$

So $\mathbb{N} \in \mathcal{A}$ and $\mathcal{A}$ consists of spaces isomorphic to their squares. Now for any $D \subset \mathbb{N}$:

$$\mathbb{N} \oplus D \cong B_0 \oplus B_1 \oplus D \cong [B_0 \cup (D \cap A_1)] \oplus [B_1 \cup (D \cap A_0)] \cong \mathbb{N} \oplus \mathbb{N} \cong \mathbb{N}$$

This in particular shows that $[e_i]_{\mathbb{N}}$ is isomorphic to its hyperplanes.

We notice now that the argument is quite general, in the sense that we could have begun from any $[e_i]_A$ instead of $[e_i]_{\mathbb{N}}$, and therefore the results hold for any space spanned by a subsequence. $\square$

Kalton [11] showed that in case an unconditional basis only has a countable number of isomorphism classes on the subsequences of the basis, then the space spanned is isomorphic to its square and hyperplanes. The above result is along the same lines and we should mention that one can get uniformity results with a bit of extra care in the proof, see the article by Ferenczi and the author, [4], for this.

Notice that permutative equivalence between subsequences of a basis induces an analytic equivalence relation on $[\mathbb{N}]^{\mathbb{N}}$.

P. Casazza drew my attention to the following theorem from [2] (proposition 6.2).

Theorem 20 (Bourgain, Casazza, Lindenstrauss, Tzafriri) *If $(e_i)_{\mathbb{N}}$ is an unconditional basic sequence permutatively equivalent to all of its subsequences, then there is a permutation π of $\mathbb{N}$ such that $(e_{\pi(i)})_{\mathbb{N}}$ is subsymmetric.*

Their statement of the theorem is slightly more general, but the general case is easily seen to follow from the infinite dimensional Ramsey theorem.

Proposition 21 *Let $(e_i)_{\mathbb{N}}$ is an unconditional basic sequence. Then either E_0 reduces to the relation of permutative equivalence of the subsequences of the basis or there is some $A \in [\mathbb{N}]^{\mathbb{N}}$ such that $(e_i)_A$ is subsymmetric.*

Proof : Notice that $\sim_p$ (permutative equivalence) on $[\mathbb{N}]^{\mathbb{N}}$ is E'_0 -invariant, so applying the Ramsey result we can suppose that there is some $B \in [\mathbb{N}]^{\mathbb{N}}$ such that all $C \in [B]^{\mathbb{N}}$ are $C \sim_p B$. Now there is some permutation π of B such that $(e_{\pi(i)})_B$ is subsymmetric. Again choosing a strictly increasing sequence $A = \{n_0, n_1, n_2, \dots\} \subset B$ such that $\pi(n_0) < \pi(n_1) < \pi(n_2) < \dots$, we get a subsymmetric $(e_i)_A$. $\square$

Acknowledgement This article was written while the author was a doctoral student in Paris under the direction of Alain Louveau. I am sincerely grateful for all his help and the interest he showed in my work.

Appendix

For the convenience of the reader we include the proof to the effect that having a minimal subspace is Σ_2^1 . This is a slightly amended version of the proof in [5] showing that having a block-minimal subspace is a Σ_2^1 property.

So suppose that X has a minimal subspace and that it does not contain c_0 . Then it has a minimal subspace with a basis (e_i) and positively separated inevitable sets F_0, F_1 in the unit sphere of $[e_i]$. (Again this is by the results of Odell and Schlumprecht.)

We let $D = \{Y = (y_i) \in bb_{\mathbb{Q}}(e_i) \mid \forall i \ y_i \in F_0 \cup F_1\}$ and as in lemma 7 let $\alpha(Y) \in 2^{\mathbb{N}}$ be defined by $\alpha(Y)(i) = 0 \leftrightarrow y_i \in F_0$. Again D is large in

$[Z]$ for any $Z \in bb_{\mathbb{Q}}(e_i)$. Take $\beta: 2^{\mathbb{N}} \longleftrightarrow (\mathbb{Q}_*^{<\mathbb{N}})^{\mathbb{N}}$ to be some fixed recursive isomorphism.

Given a $Y = (y_i) \in bb_{\mathbb{Q}}(e_i)$ any element $(\underline{\lambda}_i) \in (\mathbb{Q}_*^{<\mathbb{N}})^{\mathbb{N}}$ codes a unique infinite sequence of block vectors (not necessarily consecutive) of Y , which we denote by $(\underline{\lambda}_i) \times Y$. So due to the minimality of $[e_i]$ there is for any $Y = (y_i) \in bb_{\mathbb{Q}}(e_i)$ some $(\underline{\lambda}_i) \in (\mathbb{Q}_*^{<\mathbb{N}})^{\mathbb{N}}$ such that $(e_i) \approx (\underline{\lambda}_i) \times Y$ (a standard perturbation argument shows that the basic sequence (e_i) always embeds as a sequence of finite rational blocks, though not necessarily consecutive).

Set $W = \{Y = (y_i) \in bb_{\mathbb{Q}}(e_i) \mid (y_{2i}) \in D \wedge (e_i) \approx \beta \circ \alpha(y_{2i}) \times (y_{2i+1})\}$, which is then a Borel subset of $bb_{\mathbb{Q}}(e_i)$.

We claim that W is large in $bb_{\mathbb{Q}}(e_i)$. For suppose that $Z \in bb_{\mathbb{Q}}(e_i)$ is given, take some $V = (v_i) \in D \cap [Z]$ and a $(\underline{\lambda}_i) \in (\mathbb{Q}_*^{<\mathbb{N}})^{\mathbb{N}}$ such that $(e_i) \approx (\underline{\lambda}_i) \times (v_{3i+2})$. Choose $y_{2i} = v_{3i}$ or $y_{2i} = v_{3i+1}$ such that $\beta \circ \alpha(y_{2i}) = (\underline{\lambda}_i)$ and put $y_{2i+1} = v_{3i+2}$. Then obviously $Y \leq V \leq Z$ and $Y \in D$.

So by Gowers' theorem there is for any $\Delta > 0$ a winning strategy τ for II for producing blocks in W_{Δ} in some $Y = (y_i) \leq (e_i)$. By choosing Δ small enough and modifying τ a bit we can suppose that the vectors of even index played by II are in $F_0 \cup F_1$. So if Δ is chosen small enough, a perturbation argument shows that τ is in fact a strategy for playing blocks in W .

This shows that if X has a minimal subspace, but does not contain an isomorphic copy of c_0 , there are a basic sequence (e_i) , an element $Y \in bb_{\mathbb{Q}}(e_i)$ and a continuous function $\Theta: [Y] \rightarrow \overline{span}(e_i)^{\mathbb{N}}$ (defined by $\Theta(Z) = \beta \circ \alpha((\tau(Z))_{2i}) \times ((\tau(Z))_{2i+1})$) such that for all $Z \in [Y]$ we have $\overline{span}(\Theta(Z)) \subset \overline{span}(Z)$ and $(e_i) \approx \Theta(Z)$. On the other hand, containing a copy of c_0 is evidently a Σ_1^1 property, so the disjunction of the two becomes Σ_2^1 .

References

- [1] J. Bagaria, J. Lopez-Abad: Weakly Ramsey sets in Banach spaces. *Adv. Math.*, Vol. **160** (2001), no. 2
- [2] J. Bourgain, P. Casazza, J. Lindenstrauss, L. Tzafriri: Banach spaces with a unique unconditional basis, up to permutation. *Mem. Amer. Math. Soc. no. 54* (1985)
- [3] V. Ferenczi, E. M. Galego: Equivalence relations which are Borel reducible to isomorphism between separable Banach spaces, *preprint*.

- [4] V. Ferenczi, C. Rosendal: On the number of non isomorphic subspaces of a Banach space, *preprint*.
- [5] V. Ferenczi, C. Rosendal: Ergodic Banach spaces, *preprint*.
- [6] T. Gowers: An infinite Ramsey theorem and some Banach-space dichotomies, *Ann. of Math. 2*, Vol. **156**, (2002), no.3
- [7] T. Gowers, B. Maurey: Banach spaces with small spaces of operators, *Mathematische Annalen*, Vol. **307**, (1997)
- [8] L. Harrington, A. Kechris, A. Louveau: A Glimm-Effros dichotomy for Borel equivalence relations, *Journal of the Am. Math. Soc.*, Vol. **3**, no. 4 (1990)
- [9] L. Harrington, D. Marker, S. Shelah: Borel orderings. *Trans. Amer. Math. Soc.*, Vol. **310**, no. 1 (1988)
- [10] G. Hjorth, A. Kechris: Analytic equivalence relations and Ulm-type classifications, *Journal of symb. logic*. Vol. **60** (1995)
- [11] N. Kalton: A remark on Banach spaces isomorphic to their squares, *Contemp. Math.* **232**, Amer. Math. Soc. (1999)
- [12] A. Kechris: Classical Descriptive set theory, *Springer, New York* (1995)
- [13] R. Komorowski, N. Tomczak-Jaegermann: Banach spaces without local unconditional structure, *Israel J. Math.* **89** (1995), no. 1-3. Erratum to: "Banach spaces without local unconditional structure", *Israel J. Math.* **105** (1998).
- [14] J. Lopez-Abad: Coding into Ramsey sets, *preprint*.
- [15] E. Odell, T. Schlumprecht: The distortion problem. *Acta Math.*, Vol. **173**, no. 2.

Christian Rosendal
 Mathematics 253-37
 California Institute of Technology
 Pasadena, CA 91125
 USA
 rosendal@caltech.edu, rosendal@ccr.jussieu.fr